

28 JAN 2005

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau




(43) International Publication Date
5 February 2004 (05.02.2004)

PCT

(10) International Publication Number
WO 2004/012223 A1

- (51) International Patent Classification⁷: **H01J 49/26** (72) Inventor; and
(75) Inventor/Applicant (for US only): **KALINITCHENKO, Iouri** [AU/AU]; 5 Halifax Street, Mulgrave, Victoria 3170 (AU).
- (21) International Application Number: **PCT/AU2003/000955**
- (22) International Filing Date: **29 July 2003 (29.07.2003)** (74) Agent: **PHILLIPS ORMONDE & FITZPATRICK**; 367 Collins Street, Melbourne, Victoria 3000 (AU).
- (25) Filing Language: **English** (81) Designated States (national): **AU, CA, CN, JP, SG, US.**
- (26) Publication Language: **English** (84) Designated States (regional): **European patent (DE, FR, GB).**
- (30) Priority Data: **2002950505** **31 July 2002 (31.07.2002)** **AU** Published: **— with international search report**
- (71) Applicant (for all designated States except US): **VARIAN AUSTRALIA PTY LTD** [AU/AU]; 679 Springvale Road, Mulgrave, Victoria 3170 (AU). For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: **MASS SPECTROMETRY APPARATUS AND METHOD**



(57) Abstract: A mass spectrometer in which a substance is introduced into a plasma (28) which contains analyte ions as the plasma (28) is passing through an aperture (42), for example in a skimmer cone (40) between two vacuum regions (38) and (44) so that the substance interacts with the plasma (28) thereby reducing the concentration of interfering polyatomic or multicharged ions in the plasma by reactive or collisional interactions. The substance may be supplied via passage (60) having an outlet (63) in skimmer cone (40). The invention gives improved attenuation of interfering ions because the substance is supplied directly into the plasma (28) as it is substantially radially confined by aperture (42) and before an ion beam (58) is extracted. Alternatively or additionally a substance may be supplied directly into the plasma within aperture (36) in sampling cone (34).